

## STAB WITH KNIFE IMPACTED IN VERTEBRAL BODY AN UN-USUAL FINDING ON IMAGING

Nizamuddin Memon, Aftab Ahmed Qureshi, Muneer Junejo  
Kusum Perdeep and Abdul Sattar Memon

### ABSTRACT

Thoraco-lumbar spine injuries are caused by assault, accidents or even postural changes and are common, but this case which we report is an unusual finding in which blade of a knife was left in the vertebral body causing its fracture. A 40 years old man was brought to the emergency department with history of assault, metallic foreign body (Blade of knife) impacted in lower dorsal vertebra.

After X-ray lumbo sacral spine, CT Scan and other investigations, the injured was operated and discharged after one week. On follow-up visits, patient had no evidence of any neurological symptom/deficit.

### CASE REPORT

A 40 years old patient was brought to neurosurgery emergency department with stab injury at the back with the history of assault. Patient was fully conscious with BP 140/70 mm HG and pulse 80/minute. X-ray films showed impacted blade of a knife within T12 causing its fracture and extending to the anterior border of 12th thoracic vertebral body. The penetrated knife had partially obscured the underlying structures i.e. spinous process, lamina, pedicle and vertebral body (**Fig No. I and II**), so, computerized tomography (CT) Scan was advised for proper localization and extension of injury. CT Scan was done in prone position at Radiology Department of Liaquat University Hospital Jamshoro. Anteroposterior (AP), lateral and multiple axial scans of 10mm thickness were taken through the lower dorsal and upper lumbar spines. A metallic foreign body resembling shape of knife was seen piercing the posterior abdominal wall and splitting the muscles. There was evidence of splitting lamina and fracture of the body of 12th vertebra. Discontinuity of the cortex of the lamina of T11 vertebra was also seen along with the split of disc between T11 and T12. The interesting thing to note was that the spinal cord and

the great vessels were undamaged by distance of few millimeters. (**Fig No. III and IV**).

### OPERATIVE NOTE

The patient was operated through para spinal linear incision and diathermic cautery was done. Muscles were cut, retracted and foreign body freed. Laminectomy was done. The cord was lying by the side. It was retracted and protected. The foreign body was removed with closure done in layers.

Figure no. I and II  
showing dense metallic shadow from back to anterior border  
of 12th dorsal vertebral body





Figure no. III and IV: Metallic foreign body is seen penetrating from back to anterior border of 12th thoracic vertebral body



## DISCUSSION

We present this unusual finding on imaging which was not earlier seen in our 19 years experience. As compared with other literature, stab wounds to the spinal cord are relatively uncommon in North America but even rare in case of presentation of such injury in a delayed fashion.<sup>1</sup> Normally, the thoracic spine has slight kyphotic curve as compared with the lordotic curve of cervical and lumbar regions. These anatomic changes and difference in range of motion, increase the susceptibility to injury to thoraco-lumbar region. Nearly two thirds of thoraco-lumbar injuries occur at the levels of T12-L2.<sup>2,3</sup> Fractures are commonly the result of high energy injuries.<sup>4</sup> So, males of young age group are involved in it. In injuries with suspected

fractures of T-L spine AP and Lateral views of plain films are routinely advised, which demonstrate most acute injuries. The lateral views are most useful for evaluating alignment, vertebral compression and changes in the disc spaces.<sup>5,6,7</sup> When changes in the facet joints or parsinterarticularis are suspected, oblique views and C.T Scan are advised for proper assessment.<sup>3,8</sup> The computed tomography is necessary to define more clearly the location and extent of injuries or to clarify suggestive areas on routine radiograph. C.T Scan is most useful for evaluating patients with neurologic damage, retractor bone fragments and complex injuries before reconstruction.<sup>5,9</sup> In this case, the C.T Scan helped the surgeon in operation for removal of metallic foreign body. Neurological operative findings confirmed the imaging findings.

## REFERENCES

1. Kulkarni AV, Bhandari M, Stiver S, Reddy K. Delayed presentation of spinal stab wound. *J Emerg Med.* 2000; 18 (2):209 – 13
2. Daffner RH, Deeb ZI, Rothfus WE. Finger Prints of Vertebral Trauma –A unified Concept Based on Mechanisms. *Skeletal Radial.* 15:518-523.
3. Berquist TH. *Imaging of Orthopedic Trauma and Surgery.* WB Saunders, Philadelphia 1986
4. Stanislas MJ, Latham JM, Porter KM, Alpar EK, Stirling AJ. A High Risk Group for Thoraco Lumbar Fractures. *Injury.* 1998; 29 (1): 15-18
5. Berquist TH: *Imaging of Adult and Cervical Spine Trauma.* Radiographics. 1988; 8(4): 667-693
6. Gehweiler JA, Osborne R1, Becker Rf. *The Radiology of Vertebral Trauma.* WB Saunders, Philadelphia, 1980
7. Lauridsen KN, De Carvalho A, Anderson AH. Degree of Vertebral Wedging of the Dorsi-Lumbar Spine. *Acta Radial Diagn.* 1984; 25:29-32
8. Angtuaco EJC, Binet EF. *Radiology of Thoracic and Lumbar Fractures.* Clin Orthop. 1984; 189: 43-57
9. Acheson MB, Livingston RR, Richardson ML, Stano GK. High resolution CT Scanning in the evaluation of cervical spine trauma - A comparison with plain x-ray examination. *A J R.* 1987; 1179-85

## Correspondence:

Dr. Nizamuddin Memon  
Associate Professor, Department of Radiology & Imaging  
Liaquat University of Medical & Health Sciences  
Jamshoro, Pakistan.

