

Cupping Therapy Drains the Extra Lipids in Migraine Headache Patients

Zeba Hussain Soomro¹, Naseem Aslam Channa^{2*}, Marvi Shaikh^{2,3}, Zahoor Ahmed^{2,4},
Amna Yasin Awan², Shazia Parveen Solangi⁵

¹Department of Neurology, Civil Hospital, Karachi, Sindh-Pakistan

²Institute of Biochemistry, University of Sindh, Jamshoro, Sindh-Pakistan

³Department of Biochemistry, Indus Medical College, University of Modern Sciences, Tando Muhammad Khan, Sindh-Pakistan

⁴Institute of Public Health, Faculty of Community Medicine & Health Sciences, People's University of Medical and Health Sciences for Women, Nawabshah, Sindh-Pakistan

⁵Department of Biochemistry, Shah Abdul Latif University, Khairpur, Sindh-Pakistan

Correspondence: nachanna2000@gmail.com

doi: 10.22442/jlumhs.2025.01219

ABSTRACT

OBJECTIVES: To determine how wet cupping affects the clinical and laboratory parameters of individuals with migraine headaches.

METHODOLOGY: This cross-sectional study was conducted at the Department of Neurology, Dr Ruth K M Pfau Hospital Karachi and Hamdard University in collaboration with the Institute of Biochemistry University of Sindh Jamshoro from January 2021 to December 2023. A total of 102 migraine headache patients aged 15 to 70 years, including 32 males and 70 females, with evidence of migraine headache and normal EEG, which confirm the migraine participated in the study after providing written informed consent. Patients with abnormal EEG tests younger than 15 were excluded from the study. The patients underwent wet cupping; 5ml intravenous blood samples were collected before the procedure, 5ml from the cups, and 5ml intravenous blood was collected after one week of cupping therapy. The spectroscopic method analyzed all the samples for serum lipid profile. Data analysis was performed using (SPSS Version 26), with a significance level set as $p < 0.05$ at a 95% confidence interval.

RESULTS: Most migraine patients were female. The wet cupping samples exhibited significant increases $p < 0.05$ in lipid and liver profiles compared to pre-cupping samples, although they remained within normal ranges. No significant variations were observed in gender-specific comparisons for serum lipids. Additionally, age-wise comparisons revealed no significant differences among migraine patients of various age groups in pre-cupping and cupping samples.

CONCLUSION: Wet cupping therapy drains the extra serum lipids and reduces the migraine headache.

KEYWORDS: Migraine, Headache, Wet cupping, Intravenous, Liver function test, Lipid profile.

INTRODUCTION

World Health Organization (WHO) states that migraines impact millions of individuals globally. The NIMH (National Institute of Mental Health) states that migraines rank among the top 20 causes of disability globally¹. Cupping therapy (Al-Hijamah) has a long history in the Arabic and Muslim communities. It is believed to have been used by Prophet Muhammad (PBUH) over 1400 years ago to treat human ailments in several authentic sayings². It is a therapeutic technique that involves using suction, usually through a fire-generated vacuum cup or vessel, on the affected area or any surface of the body to treat disease³. Hijamah in Arabic is the process of applying cups to migraine. Headache is one of the most prevalent human disorders worldwide⁴. Cupping does not function to eliminate blood but to extract metabolic waste known as causative pathological substances⁵. It is usually used to remove blood and fluid from abnormally elevated causes of pathological conditions, such as high lipid serum levels⁶. Some of the studies indicated that wet cupping can reduce pain. A limited number of studies have been conducted to evaluate the effectiveness of wet cupping in treating migraine headaches⁷. The association between migraine and associated vascular and liver diseases has long been established. The contribution of these risk factors to this association remains unclear⁸. Therefore, there is great potential if wet cupping therapy can be used to reduce these disease risk factors such as lipid profile and Liver function. Thus, this study explored the role of wet cupping therapy on lipid profile and liver function test in migraine headache patients.

METHODOLOGY

This descriptive cross-sectional study employed non-probability consecutive sampling. This study was conducted at the Department of Neurology, Dr. Ruth K. M. Pfau, Civil Hospital, Karachi and Hamdard University in collaboration with the Institute of Biochemistry University of Sindh Jamshoro from January 2021 to December 2023. One hundred and two patients diagnosed with migraine headaches were included. Patients with normal EEG confirming the migraine participated in the study after providing written informed consent were included. Patients with abnormal EEG tests younger than 15 years were excluded from the study. Ethical approval was obtained from the Bioethical Committee via letter no. IOB/53/2019, dated 18/1/2019. A questionnaire was administered to collect the socio-demographic characteristics of the migraine patients. Participants were asked to fast for at least 3 hours before the procedure. Approximately 5 ml of intravenous blood was collected five minutes before the wet cupping therapy procedure in a serum-separating tube, and an additional 5 ml blood sample was collected using 5 cc disposable syringes from the cupping site. After one week of cupping Therapy, another 5 ml of venous blood was collected. Before cupping, wet cupping and after-wet cupping samples were compared for selected biochemical parameters. Intravenous and wet cupping blood samples were centrifuged at 3000 rpm for 15 minutes, and the serum was stored at -68°C for lipid analysis using the Lab L19 ELISA 6000 Cobas Roche Analyzer, following the rules of the International Federation of Clinical Chemistry (IFCC).

Statistical analysis: Statistical analysis was performed using SPSS Version 26, with a p-value < 0.05 considered significant and a 95% confidence interval applied.

RESULTS

Table I: General socio-demographic characteristics of the migraine headache patients

Characteristics	Frequency (n=102)	Percentage %
Gender		
Male	32	31.4
Female	70	68.6
Mean age (year)	35.18 ± 12.63	
Education		
Illiterate	28	27.50
Primary	30	29.40
Secondary	13	12.70
Higher	31	30.40
Socioeconomic status		
Lower class	79	77.50
Middle class	10	9.80
Upper class	13	12.70

Table I illustrates the general socio-demographic characteristics of migraine headache patients. Among the 120 participants, the majority (68.60%) were females, with a mean age of 35.18. Most of the migraine headache patients were found to be literate, followed by illiterate patients. The socioeconomic status of the migraine headache patients was primarily low, followed by middle and high classes.

Table II: Comparison of biochemical parameters in before, wet cupping and after cupping samples of migraine headache patients

Biochemical parameters	Sample type	Migraine headache patients n=102 Mean ± SD	Male n=32 Mean ± SD	Female n=70 Mean ± SD
SERUM LIPIDS				
Total Cholesterol < 200 (mg/dL)	BCS	161.66±23.33	171.20±22.71 ^b	170.30±22.20 ^c
	WCS	170.62±22.30 ^a	192.88±41.04 ^b	182.63±41.04 ^c
	ACS	169.33±69.80 [*]	148.50±45.43	178.86±76.14
Triglycerides < 150 (mg/dL)	BCS	117.20±23.48	129.10±21.32 ^b	160.79±85.66 ^c
	WCS	126.65±22.12 ^a	123.00±21.41 ^b	174.73±81.19 ^c
	ACS	150.81±141.7	121.28±121.8	164.31±146.9
High-Density Lipoproteins < 40 (mg/dL)	BCS	40.97±3.690	44.41±3.670 ^b	44.01±4.690 ^c
	WCS	44.13±4.366 ^a	41.48±2.553 ^b	40.75±4.110 ^c
	ACS	42.92±18.80 [*]	38.55±11.77	44.94±20.83
Low-Density Lipoproteins < 100 (mg/dL)	BCS	81.99±9.414	96.81±29.13 ^b	85.91±10.33 ^c
	WCS	85.82±9.293 ^a	107.75±35.04 ^b	82.01±10.52 ^c
	ACS	95.28±45.97	85.88±30.89	99.59±50.54
Very Low-Density Lipoproteins <30 (mg/dL)	BCS	28.048±11.26	35.9±12.92 ^b	35.21±13.68 ^c
	WCS	35.43±13.36 ^a	40.57±26.51 ^b	28.04±11.40 ^c
	ACS	30.16±28.34 [*]	24.26±24.37	32.86±29.39
Total lipids 450-1000 (mg/dL)	BCS	170.12±34.08	170.76±32.58	169.83±34.70 ^c
	WCS	171.86±61.03 ^a	192.83±41.00	162.27±65.66 ^c
	ACS	168.39±67.41	148.68±40.26	177.40±74.54
LFTs				
Alkaline phosphatase M= 50-136; F= 45-136 (U/L)	BCS	86.36±23.54	87.65±24.60	92.75±23.51 ^c
	WCS	91.08±23.87 ^a	85.75±27.00	86.64±21.93 ^c
	ACS	97.14±72.11	91.00±50.83	99.94±79.34
Alanine transmarine M=50 (U/L); F= 35 (U/L)	BCS	17.19±7.719	21.61±9.00 ^{ab}	17.53±7.470 ^{ac}
	WCS	18.86±8.190 ^a	18.84±8.920 ^b	16.43±7.032 ^c
	ACS	19.46±17.61	21.47±18.99	18.54±16.73
Bilirubin < 1.2 (mg/dL)	BCS	0.332±0.140	0.434±0.140	0.340±0.153 ^c
	WCS	0.370±0.148 ^a	0.450±0.12 ^a	0.360±0.140 ^{ac}
	ACS	0.360±0.284	0.300±0.169	0.390±0.318


a=p <0.05 for the comparison between intravenous and cupping samples. *= p <0.05 for comparing cupping and after-cupping samples. b=p <0.05 for comparing male patients between intravenous and cupping samples. c=p <0.05 for the comparison of female patients in before cupping and cupping samples, ACS= After cupping sample, BCS, Before cupping sample, WCS= Wet cupping sample

In the present study, the comparison of serum lipids and liver function tests in intravenous and wet cupping and after-cupping samples are presented in **Table II**. The total cholesterol concentration in wet cupping serum significantly exceeded that in the before-cupping samples. The low-density lipoprotein (LDL), high-density lipoprotein (HDL), triglycerides (TG), very low-density lipoprotein (VLDL), and total lipids in the wet cupping samples were significantly

higher compared to the before cupping samples. Both (HDL and LDL) were also considerably higher in the cupping samples when compared to the after-cupping samples. In this study, serum (ALT), (ALP) and bilirubin concentrations in wet-cupping samples were significantly higher compared to the before-cupping samples. No significant variation was found when comparing before and after cupping samples and cupping samples with after cupping samples.

A statistically significant increase in cholesterol, triglycerides and (LDL), (VLDL) and (HDL) levels was observed in the wet cupping samples of females compared to intravenous samples (**Table III**). As indicated in **Table III**, there were no noteworthy variations in lipid profile when comparing intravenous and wet cupping with after-cupping samples across genders. Among male patients, (ALP) and bilirubin levels were significantly increased in intravenous samples compared to wet cupping samples. No significant changes in (ALP) and bilirubin were observed in the male patients in either group.

All serum lipids increased significantly in the wet cupping samples compared to the intravenous samples in males with migraine headaches. The current study found that cupping Therapy drains cholesterol and (LDL) in male migraine headache patients.



ACS= After cupping sample, BCS, Before cupping sample, WCS= Wet cupping sample, TC=Total cholesterol, TG= Triglycerides, HDL= High density lipoproteins, LDL= Low density lipoproteins, VLDL=Ver low density lipoproteins, TL= Total lipids

Figure I: Age-wise comparison of lipid profile in samples of migraine headache patients before cupping, wet cupping, and after cupping.

Table III: Age-wise comparison of liver function tests in samples of migraine headache patients before cupping, wet cupping, and after cupping

LFTs	Sample type	≤ 15 - 30 y	31 - 45 y	46 - 60 y	≥ 61 y	P-value > 0.05
		Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	
ALP (U/L) M= 50-136 F= 45-136	BCS	100.84 ± 3.690	85.838 ± 4.719	93.904 ± 7.690	124.0 ± 19.0	0.071
	WCS	105.14 ± 5.197	92.428 ± 4.845	98.666 ± 8.939	94.01 ± 2.80	0.651
	ACS	105.70 ± 9.194	88.19 ± 5.143	89.141 ± 3.309	154.00 ± 6.60	0.021
ALT (U/L) M=50 F= 35	BCS	18.913 ± 6.271	20.903 ± 8.288	20.857 ± 3.188	17.00 ± 2.01	0.213
	WCS	21.642 ± 8.305	22.464 ± 7.861	21.380 ± 3.712	14.50 ± 5.50	0.120
	ACS	18.07 ± 5.903	21.520 ± 2.046	20.330 ± 3.929	19.50 ± 2.500	0.341
BILIRUBIN < 1.2 (mg/dL)	BCS	0.3826 ± 0.256	0.3441 ± 0.168	0.375 ± 0.261	0.375 ± 0.035	0.062
	WCS	0.4385 ± 0.266	0.3507 ± 0.148	0.4109 ± 0.254	0.320 ± 0.091	0.154
	ACS	0.3620 ± 0.281	0.3701 ± 0.301	0.3410 ± 0.278	0.431 ± 0.160	0.810

ACS= After cupping sample, BCS, Before cupping sample, WCS= Wet cupping sample, ALP= Alkaline phosphatase, ALT= Alanine transferase

There were no notable differences in serum lipid levels among the age groups of migraine headache patients in samples collected before cupping, during cupping, and after wet cupping, except for HDL. HDL levels had been significantly lower in samples collected after one week of wet cupping therapy across all age groups. Additionally, as shown in **Table III**, there had been a significant increase in ALP concentrations in the age group of >61 years and wet cupping samples compared to other age groups. When comparing the same age group in intravenous and post-wet cupping samples of migraine headache patients, the wet cupping samples showed a significant decrease in ALP levels. This emphasized the significance of wet cupping therapy, particularly in the elderly population.

DISCUSSION

In **Table I**, women are more likely to suffer from migraine, but this may be due to pathophysiology influenced by sexual hormones. For example, Progesterone and testosterone have not been studied as extensively in migraine patients and estrogen withdrawal is known to cause migraine⁹. According to a US Migraine Prevalence and Prevention Study (AMPP), 43% of women and 18% of men suffered from migraines during their lifetime¹⁰. These results also corroborate the present study, as we found that men were more numerous than women. Studies on the association between socio-demographic status and migraine headaches have yielded mixed results. Several studies have shown that migraine incidence is more common among lower-income and low-education groups, which is similar to the findings of this study. The low social and economic status of education and occupation is related to an increased risk of frequent and chronic headaches¹¹.

The research results shown in **Table II** demonstrated that wet cupping therapy significantly affected total cholesterol, TG, HDL, LDL, and VLDL. Previous studies revealed a reduction in lipid profile by sampling intravenous blood before and after wet cupping therapy; the wet cupping therapy blood was laden with elevated lipid profiles, resulting in a reduction in intravenous blood lipid profiles after wet cupping therapy¹². The effect of wet cupping therapy on blood detoxification could also be supported by a previous study that compared wet cupping blood cholesterol levels with venous blood cholesterol levels. The results showed that wet cupping therapy blood cholesterol was higher than intravenous blood¹³. The same findings were also reported in the present study. Wet cupping could potentially serve as a practical approach to lower LDL cholesterol levels in men, thereby potentially offering a preventive impact against the development of atherosclerosis¹⁴. Numerous physicians assert that cupping is a preventive measure against certain cardiovascular diseases¹⁵.

High serum cholesterol level is a significant risk factor for the development of cardiovascular disease. It is known that cholesterol levels in the expelled blood are significantly higher during cupping than in the intravenous blood^{16,17}. Triglycerides, HDL, cholesterol, LDL, and VLDL concentrations show variations between the three groups, although none of the differences reached statistical significance (p -values < 0.05), as shown in **Table III**. This suggests that cupping Therapy did not significantly affect the lipid profile in female migraine headache patients¹⁸. Previous studies have also demonstrated that cupping blood is more detoxifying than intravenous blood¹⁹.

CONCLUSION

Cupping contributes to the improvement of blood components in individuals. It can be presumed that a distinct variation exists in the composition of blood obtained from cupping in contrast to the blood collected intravenously before and following Therapy. Cupping Therapy helps to improve serum cholesterol levels for migraine headache patients by reducing low-density lipoprotein (LDL) and total cholesterol levels. It also helps to lower some biochemical parameters that can be associated with metabolic diseases. This means that cupping therapy may help treat many diseases such as heart and liver diseases by decreasing the body's components.

HUMAN ETHICAL APPROVAL AND CONSENT TO PARTICIPATE

Ethical approval for this study was obtained from the Ethical Review Committee of the University of Sindh Jamshoro. The Institutional Ethical Committee via letter no. IOB/53/2019, dated 18/1/2019, approved the study before researching migraine headache patients. Written informed consent was obtained from each participant after explaining the study's purpose and procedure. The research was conducted with complete transparency, and every possible safety measure was taken to protect all participants from potential harm or contamination of the privacy and confidentiality of the study respondents. The blood samples of all participants were collected under the supervision of a competent and licensed healthcare professional.

ACKNOWLEDGEMENTS

We sincerely thank the administration of the Department of Neurology, Dr. Ruth K. M. Pfau Civil Hospital Karachi, for providing moral support for the success of our research.

Ethical permission: University of Sindh, IRB letter No. IOB/53/2019.

Conflict of Interest: No conflicts of interest, as stated by authors.

Financial Disclosure / Grant Approval: The authors did not receive any funding from any source, and the research was carried out through facilities provided by the Institute of Biochemistry, University of Sindh, Jamshoro.

Data Sharing Statement: The corresponding author can provide the data proving the findings of this study on request. Privacy or ethical restrictions bound us from sharing the data publicly.

AUTHOR CONTRIBUTION

Soomro ZH: Responsible for the conception and design of the study

Channa NA: Responsible for the conception and design of the study, data interpretation, and substantial revisions to enhance the quality of the final work.

Shaikh M: Data collection, blood sampling, experimental analysis, prepared the initial draft of the manuscript, provided expertise in the methodology and supported the editing and proofreading.

Ahmed Z: Data collection, blood sampling, and experimental analysis.

Awan AY: Data collection, blood sampling, experimental analysis, assisted with the graphical representation of data and the preparation of figures and tables.

Solangi SP: Data collection, blood sampling, and experimental analysis.

REFERENCES

1. Shan C-S, Xu Q-Q, Shi Y-H, Wang Y, He Z-X, Zheng G-Q. Chuanxiong formulae for migraine: a systematic review and meta-analysis of high-quality randomized controlled trials. *Frontiers in Pharmacology*. 2018;9:589.
2. Qureshi NA, Ali GI, Abushanab TS, El-Olemy AT, Alqaed MS, El-Subai IS, et al. History of cupping (Hijama): a narrative review of literature. *Journal of integrative medicine*. 2017;15(3):172-81.
3. Al Bedah AM, Khalil MK, Posadzki P, Sohaibani I, Aboushanab TS, AlQaed M, et al. Evaluation of wet cupping therapy: systematic review of randomized clinical trials. *The Journal of Alternative and Complementary Medicine*. 2016;22(10):768-77.
4. Aboushanab TS, AlSanad SM. Cupping therapy (Hijama) in the Arab world. *Handbook of Healthcare in the Arab World*: Springer; 2021. p. 1845-64.
5. El Sayed SM, Mahmoud HS, Nabo MMH. Methods of wet cupping therapy (Al-Hijamah): in light of modern medicine and prophetic medicine. *Altern Integ Med*. 2019;2(3):1-16.
6. El-Shanshory M, Hablas NM, Shebl Y, Fakhreldin AR, Attia M, Almaramhy HH, et al. Al-hijamah (wet cupping Therapy of prophetic medicine) significantly and safely reduces iron overload and oxidative stress in thalassemic children: a novel pilot study. *Journal of blood medicine*. 2018;9:241-51.
7. Aboushanab TS, AlSanad S. Cupping therapy: an overview from a modern medicine perspective. *Journal of acupuncture and meridian studies*. 2018;11(3):83-7.
8. Liampas I, Mylonas KS, Brotis A, Dervenis P, Siokas V, Mentis AFA, et al. Serum lipid abnormalities in migraine: A meta-analysis of observational studies. *Headache: The Journal of Head and Face Pain*. 2021;61(1):44-59.
9. Ahmad SR, Rosendale N. Sex and gender considerations in episodic migraine. *Current pain and headache reports*. 2022;26(7):505-16.
10. Allais G, Chiarle G, Sinigaglia S, Benedetto C. Menstrual migraine: a review of current and developing pharmacotherapies for women. *Expert opinion on pharmacotherapy*. 2018;19(2):123-36.
11. Westergaard ML, Glümer C, Hansen EH, Jensen RH. Prevalence of chronic headache with and without medication overuse: associations with socioeconomic position and physical and mental health status. *PAIN®*. 2019;155(10):2005-13.
12. Allafi AR, Al-Haifi AR. The effect of Hijamah on different health parameters. *PROGRESS IN NUTRITION*. 2020;22(2):411-4.
13. Saeed AA, Badulla WF, Sheikh GAA. The effect of wet cupping therapy (al-hijamah) on some blood components: a comparative study. *Electronic Journal of University of Aden for Basic and Applied Sciences*. 2021;2(3):124-30.
14. Niasari M, Kosari F, Ahmadi A. The effect of wet cupping on serum lipid concentrations of clinically healthy young men: a randomized controlled trial. *The Journal of Alternative and Complementary Medicine*. 2017;13(1):79-82.
15. Shekarforoush S, Foadoddini M, Noroozadeh A, Akbarinia H, Khoshbaten A. Cardiac effects of cupping: myocardial infarction, arrhythmias, heart rate and mean arterial blood pressure in the rat heart. *Chin J Physiol*. 2017;55(4):253-8.
16. Wadhera RK, Steen DL, Khan I, Giugliano RP, Foody JM. A review of low-density lipoprotein cholesterol, treatment strategies, and its impact on cardiovascular disease morbidity and mortality. *Journal of Clinical Lipidology*. 2016;10(3):472-89.
17. Lu S, Du S, Fish A, Tang C, Lou Q, Zhang X. Wet cupping for hypertension: a systematic review and meta-analysis. *Clinical and Experimental Hypertension*. 2019;41(5):474-80.
18. Shekarforoush S, Foadoddini M, Noroozadeh A, Akbarinia H, Khoshbaten A. Cardiac effects of cupping: myocardial infarction, arrhythmias, heart rate and mean arterial blood pressure in the rat heart. *Chinese Journal of Physiology*. 2018;55(4):253-8.
19. Al-Bedah AM, Elsubai IS, Qureshi NA, Aboushanab TS, Ali GI, El-Olemy AT, et al. The medical perspective of cupping therapy: Effects and mechanisms of action. *Journal of traditional and complementary medicine*. 2019;9(2):90-7.