

Diagnostic Accuracy of Ripasa and Alvarado Score in Detecting Acute Appendicitis

Shahid Nazir Memon^{1*}, Shahnawaz Abro¹, Muhammad Rafique Pathan¹, Anita Kumari¹,
Rehmat-Urf-Sehrish¹, Sahrish Sulman¹

ABSTRACT

OBJECTIVE: To determine RIPASA and ALVARADO score diagnostic accuracy in detecting acute appendicitis.

METHODOLOGY: A cross-sectional comparative study was performed on patients aged ≥ 18 at the Liaquat University of Medical and Health Sciences Jamshoro Surgery Department. Two hundred and thirty patients were evaluated from April to September 2019. Patients with right iliac fossa (RIF) pain were consecutively selected. Patients with right iliac fossa mass, pregnant women, having pelvic inflammatory disease, or urolithiasis history were excluded from the study. RIPASA and ALVARADO scores were applied to each patient, and histopathology was used as the gold standard. Data analysis was performed with SPSS version 25.

RESULTS: Male were 132 (57.4%) and female were 98 (42.6%) presented with suspected acute appendicitis with a mean age of 29.8 ± 9.2 years. Histopathology confirmed acute appendicitis in 217 (94.3%) patients, whereas RIPASA score in 210 (91.3%) patients and ALVARADO score in only 30 (13.0%) patients. Sensitivity (94.01% vs 13.36%), specificity (53.85% vs 92.31%), positive predictive value (PPV, 97.14% vs 96.67%), negative predictive value (NPV, 35.00% vs 6.00%) and diagnostic accuracy (DA, 91.74% vs 17.83%) of RIPASA and ALVARADO respectively.

CONCLUSION: RIPASA score is more sensitive and accurate in predicting acute appendicitis than the ALVARADO score.

KEYWORDS: Acute appendicitis, abdominal, emergency, pain, RIPASA, ALVARADO.

INTRODUCTION

Acute appendicitis is the most frequent reason for abdominal surgery in an emergency, with a 7.0-8.0% lifetime risk¹. The reported incidence of acute appendicitis is 233 per 0.1 million, with a mean age of 28 years. Males are more commonly affected with acute appendicitis, with a lifetime risk of 8.6%, than females with 6.7%². Despite acute appendicitis being the most common health problem, its diagnosis is still difficult, specifically in children, young adults, and pregnant women. Early and accurate diagnosis is essential for decreasing perforation and worse appendectomy outcomes. Most emergency patients have signs and symptoms similar to diseases like genitourinary or gynecologic inflammatory conditions. Therefore, diagnosis of acute appendicitis depends upon presenting complaints, physical examination, laboratory investigations, and different predicting scoring systems³⁻⁵.

Different simple, inexpensive, non-invasive and reliable scoring systems are developed to diagnose acute appendicitis rapidly^{5,6}. These systems decrease the negative appendectomy rate by approximately 5-

10%. Alvarado score is the oldest and most widely used, with clinically approved diagnostic accuracy. It is the most commonly used and popular score among surgeons because it is simple, inexpensive and quick. It comprises eight predictive clinical factors with a maximum of ten score with a cut-off of ≥ 7 score^{7,8}. The Raja Isteri Pengiran Anak Saleha Appendicitis (RIPASA) score is among the most widely used scoring systems worldwide. It comprises 15 predictive parameters with a maximum of 16 score with a cut-off of ≥ 7.5 . RIPASA score has a few additional parameters for predicting acute appendicitis, which is absent in the Alvarado score of^{9,10}.

Most of these scoring systems are developed for the population of Western countries; for example, initially, the Alvarado score did not apply to Asian countries. Differences in ethnicity and diet can change the predictability of different scoring systems. Therefore, these scoring systems should be applied to the local population for more accurate results in predicting acute appendicitis. The study focuses on finding RIPASA and ALVARADO score diagnostic accuracy in detecting acute appendicitis.

METHODOLOGY

A cross-sectional comparative study was performed at the Department of Surgery, Liaquat University of Medical and Health Sciences Jamshoro, after obtaining approval from the research ethics committee (LUMHS/REC/-781). During the study period from

¹Department of Surgery, Liaquat University of Medical & Health Sciences (LUMHS), Jamshoro, Sindh-Pakistan.

*Correspondence: shahidnazir2702003@yahoo.com

doi.10.22442/jlumhs.2022.00912

Received: 25-08-2021

Revised: 21-02-2022

Accepted: 13-04-2022

Published Online: 12-05-2022



April to September 2019, 230 patients aged ≥ 18 years who presented with right iliac fossa (RIF) pain were consecutively selected. Patients without RIF pain, right iliac fossa mass, pregnant women, pelvic inflammatory disease or urolithiasis history were excluded from the study.

Suspected acute appendicitis was defined in the right iliac fossa pain presence. RIPASA score consists of 15 parameters with a total score of 16 and a cut-off of ≥ 7.5 . ALVARADO score consists of 08 parameters with a score of 10. ALVARADO score of ≥ 7.0 was used to confirm acute appendicitis (**Table I**).

The detailed medical history of each patient was obtained. RIPASA and ALVARADO scores were applied to each patient, and histopathology was used as the gold standard for confirming both scores' diagnoses. Data was interpreted using the Statistical Package for Social Science (SPSS) version 25.

TABLE I:
RIPASA AND ALVARADO SCORE CRITERIA

RIPASA Score		ALVARADO Score	
Characteristics	Score	Characteristics	Score
Gender			
Male	1.0	--	--
Female	0.5	--	--
Age			
≤ 40	1.0	--	--
> 40	0.5	--	--
Symptoms			
RIF pain	0.5	Migration of pain to the right lower quadrant	1.0
Pain migration to right iliac fossa (RIF)	0.5	Nausea & Vomiting	1.0
Anorexia	1.0	Anorexia	1.0
Nausea & Vomiting	1.0	--	--
Duration of symptoms			
≤ 48 Hours	1.0	--	--
> 48 Hours	0.5	--	--
Signs			
RIF tenderness	1.0	Tenderness in RIF	2.0
Guarding	2.0	Rebound tenderness in RIF	1.0
Rebound tenderness	1.0	Elevated temperature (> 37.3 °C)	1.0
Rovsing's Sign	2.0	--	--
Temperature: >37 °C & <39 °C	1.0	--	--
Labs			
Raised WBC count	1.0	Leukocytosis > 10000 mm ³	2.0
Negative urine analysis (Absence of blood, WBCs, bacteria)	1.0	Shift to the left of Neutrophils	1.0
Additional Scores			
Foreign I.C.	1.0	--	--
Total Score	16		10

RESULTS

Out of 230 suspected patients, males were 132 (57.4%), and females were 98 (42.6%). 139 (60.4%) patients fall in the age group of 18-30 years, and 66 (28.7%) patients in the age group of 31-40 years with a mean age of 29.8 ± 9.2 years (**Table II**).

Out of 230 suspected patients, histopathology confirmed acute appendicitis in 217 (94.3%) patients, whereas RIPASA score in 210 (91.3%) patients and ALVARADO score in only 30 (13.0%) patients. The mean RIPASA score was 11.0 ± 2.7 , and the ALVARADO score was 5.4 ± 1.4 (**Table II**).

On RIPASA score, 210 (91.3%) patients were suspected of acute appendicitis, among which histopathology (true positive) confirmed acute appendicitis in 204 (94.0%) patients, 7 (53.8%) were true negative, 6 (46.2%) were falsely positive, and 13 (6.9%) were false negative. Similarly, on the ALVARADO score, only 30 (13.0%) patients were suspected of acute appendicitis, among which histopathology (true positive) confirmed acute appendicitis in 29 (13.4%) patients, 12 (92.3%) were true negative, 1 (7.7%) was false positive and 188 (86.6%) were false negative (**Table III**).

The Sensitivity recorded (94.01% vs 13.36%), specificity (53.85% vs 92.31%), positive predictive value (PPV, 97.14% vs 96.67%), negative predictive value (NPV, 35.00% vs 6.00%) and diagnostic accuracy (DA, 91.74% vs 17.83%) of RIPASA and ALVARADO respectively (**Table III**). The receiver operating characteristic (ROC) curve also indicates a higher DA of RIPASA than Alvarado. The area under the curve (AUC) of RIPASA and Alvarado scores was 0.761 and 0.531, respectively (**Figure I**).

TABLE II: DEMOGRAPHICS AND DIAGNOSIS OF ACUTE APPENDICITIS PATIENTS (n=230)

Variables	Frequency	Percentage
Gender		
Male	132	57.4
Female	98	42.6
Age		
Mean \pm SD	29.8 ± 9.2 (18-58)	
18-30	139	60.4
31-40	66	28.7
41-50	16	7.0
51-60	9	3.9
Histopathology		
+VE	217	94.3
-VE	13	5.7
RIPASA Score		
Mean \pm SD	11.0 ± 2.7 (5.0-16.0)	
+VE	210	91.3
-VE	20	8.7
ALVARADO Score		
Mean \pm SD	5.4 ± 1.4 (4-10)	
+VE	30	13.0
-VE	200	87.0

**FIGURE I:
ROC CURVE OF RIPASA AND ALVARADO SCORE**




TABLE III: DIAGNOSTIC FINDINGS AND SENSITIVITY ANALYSIS (n=230)

Variables	Histopathology (Gold Standard)		P-value
	Present (n=217)	Negative (n=13)	
RIPASA Score			
+VE	204 (94.0%)	6 (46.2%)	< 0.001*
-VE	13 (6.9%)	7 (53.8%)	
ALVARADO Score			
+VE	29 (13.4%)	1 (7.7%)	0.55
-VE	188 (86.6%)	12(92.3%)	
Sensitivity Analysis			
	RIPASA	ALVARADO	
Sensitivity	94.01%	13.36%	
Specificity	53.85%	92.31%	
Positive Predictive Value	97.14%	96.67%	
Negative Predictive Value	35.00%	6.00%	
Diagnostic Accuracy	91.74%	17.83%	

*P-values are calculated on the Chi-square test and significant on ≤ 0.05

DISCUSSION

Acute appendicitis diagnosis is very challenging for emergency physicians. It requires a detailed patient's medical history, presenting clinical signs and symptoms, and laboratory investigations, but it is still considered a diagnostic dilemma. Therefore, various diagnostic scoring systems are utilized for the early diagnosis of acute appendicitis. The current study evaluated 230 suspected patients of acute appendicitis using RIPASA and ALVARADO scores followed by histopathology to confirm their predicted diagnosis.

Most patients with RIF pain were males in the current study; 132 (57.4%) and 98 (42.6%) were females. Most patients were young adults with a mean age of

29.8±9.2 years. Similar Pakistani studies, such as Damani SAAR 2016¹¹ and Davis GN 2019⁹, also reported a higher male prevalence of 64.5% and 58.4%, with a mean age of 24.7±10.2 and 23.5±9.1 years. Similar to studies such as Shuaib A et al.¹², Özsoy Z 2017¹³, Chisthi MM 2020¹⁴ and Kothari D 2017¹⁵ also reported a higher male prevalence mean age. All studies help conclude that young male adults are more prone to developing acute appendicitis.

In the current study, the gold standard diagnostic of acute appendicitis, i.e., histopathology, confirmed acute appendicitis in 217 (94.3%) patients. In contrast, the RIPASA score was also accurate and diagnosed acute appendicitis in 210 (91.3%) patients, but the ALVARADO score was inappropriate and diagnosed acute appendicitis in only 30 (13.0%) patients. Damani SAAR 2016¹¹ diagnosed acute appendicitis in 90.0% of patients based on histopathology, 91.1% of patients based on RIPASA score and 11.7% of patients based on ALVARADO score. Histopathology is the gold standard in diagnosing acute appendicitis, while the results of the RIPASA score are almost similar to histopathology.

In the current study, the diagnostic accuracy of the RIPASA score was much higher than the ALVARADO score. Sensitivity (94.01% vs 13.36%), specificity (53.85% vs 92.31%), positive predictive value (97.14% vs 96.67%), negative predictive value (35.00% vs 6.00%) and diagnostic accuracy (91.74% vs 17.83%) of RIPASA and ALVARADO respectively. Damani SAAR 2016¹¹ reported similar results with higher diagnostic accuracy of the RIPASA score than the ALVARADO score. Sensitivity (91.11% vs 11.67%), specificity (60.0% vs 95.0%), PPV (95.34% vs 95.45%), NPV (42.85% vs 10.67%) and DA (88.0% vs 20.0%) of RIPASA and ALVARADO score respectively. Davis GN 2019⁹ also reported the higher Sensitivity (96.7%), specificity (93.0%), PPV (94.8%), NPV (95.54%) and DA (95.1%) of RIPASA score. Studies^{12,14,16-21} also reported similar results, such as the RIPASA score being more appropriate with higher Sensitivity, specificity, and diagnostic accuracy.

In this study, the ROC curve's area under the curve also showed a higher diagnostic value of RIPASA score than ALVARADO, with higher AUC of 0.761 and 0.531, respectively. Different other studies also reported a higher AUC with RIPASA than ALVARADO to score, such as Damani SAAR 2016¹¹ 0.889 vs 0.633, Chisthi MM 2020¹⁴ 0.910 vs 0.726, Pasumarthi V 2018¹⁶ 0.810 vs 0.771 and Sanjive JG 2019¹⁷ 0.920 vs 0.490 in RIPASA and ALVARADO score respectively as ROC curve shows that RIPASA score has higher diagnostic value and AUC than ALVARADO score. It helps to conclude that the RIPASA scoring system is a more powerful diagnostic tool than ALVARADO in diagnosing acute appendicitis.

However, no any diagnostic scoring system or laboratory investigation is 100.0% reliable in diagnosing acute appendicitis. Our study findings

conclude that the accuracy of diagnosing acute appendicitis is directly associated with a higher score of RIPASA and ALVARADO. RIPASA scoring system is more reliable than ALVARADO and can be used to decide whether to proceed with patient surgery or conservative treatment. Suspected acute appendicitis patients with a score of 7.5 or greater should be managed with surgery. Patients with a score of 7 or less should be handled with conservative treatment and kept under continuous observation. If the score exceeds 7.5 or greater, the patient will undergo surgery. If the score falls below 5, the patient will be discharged from the hospital and advised to come to the hospital immediately if symptoms persist or worsen.

CONCLUSION

RIPASA score is a more sensitive, reliable, and accurate diagnostic tool for predicting acute appendicitis than the ALVARADO score. RIPASA scoring system uses simple parameters that help the surgeons reach the final decision in patients with RIF pain in an emergency.

ABBREVIATIONS

AUC: Area Under the Curve

DA: Diagnostic Accuracy

NPV: Negative Predictive Value

PPV: Positive Predictive Value

RIF: Right Iliac Fossa

RIPASA: Raja Isteri Pengiran Anak Saleha Appendicitis

ROC: Receiver Operating Characteristic

SPSS: Statistical Package for Social Science

Ethical Permission: Liaquat University of Medical & Health Sciences REC letter No. LUMHS/REC/-781.

Conflicts of Interest: The authors have no conflicts of interest to declare.

Financial Disclosure / Grant Approval: No funding agency was used for this research.

Data Sharing Statement: The data supporting this study's findings are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

AUTHOR CONTRIBUTION

Nazir S: Main Idea and Design of Study

Abro S: Collection of data and analysis

Pathan MR: Interpretation of data

Kumari A: Compilation and finalizing of the manuscript

Sehrish R: Critical review of the manuscript

Sulman S: Reference collection

REFERENCES

- Sartelli M, Baiocchi GL, Di Saverio S, Ferrara F, Labricciosa FM, Ansaloni L et al. Prospective observational study on acute appendicitis worldwide (POSAW). *World J Emerg Surg.* 2018; 13: 19. doi: 10.1186/s13017-018-0179-0.
- Jones MW, Lopez RA, Deppen JG. Appendicitis. [Updated 2021 Feb 8]. In: StatPearls [Internet]. Treasure Island (F.L.): StatPearls Publishing; 2021 Jan.
- Eng KA, Abadeh A, Ligocki C, Lee YK, Moineddin R, Adams-Webber T et al. Acute appendicitis: a meta-analysis of the diagnostic accuracy of U.S., CT, and MRI as second-line imaging tests after an initial US. *Radiology.* 2018; 288(3): 717-27. doi: 10.1148/radiol.2018180318.
- Alvarado A. How to improve the clinical diagnosis of acute appendicitis in resource limited settings. *World J Emerg Surg.* 2016; 11: 16. doi: 10.1186/s13017-016-0071-8.
- Alvarado A. Diagnostic scores in acute appendicitis. *Curr Issues Diag Treat Acute Appendicitis.* 2018; 43. doi: 10.5772/intechopen.77230.
- Xingye W, Yuqiang L, Rong W, Hongyu Z. Evaluation of diagnostic scores for acute appendicitis. *J Coll Physicians Surg Pak.* 2018; 28(2): 110-4. doi: 10.29271/jcpsp.2018.02.110.
- Tekeli MT, Ilhan E, Ureyen O, Senlikci A, Yeldan E, Ozturk M et al. How much reliable is Alvarado scoring system in reducing negative appendectomy? *Indian J Surg.* 2017; 79(2): 106-10. doi: 10.1007/s12262-015-1433-2.
- Al-Awaysih MM, Nofal MN, Yousef AJ. Evaluation of Alvarado score in diagnosing acute appendicitis. *Pan Afr Med J.* 2019; 34: 15. doi: 10.11604/pamj.2019.34.15.17803.
- Davis GN, Swaminathan M. Acute appendicitis - new diagnostic algorithm using RIPASA score and non-contrast computed tomography scan. *J Pak Med Assoc.* 2019; 69(Suppl 1)(1): S12-S16.
- Malik MU, Connelly TM, Awan F, Pretorius F, Fiuza-Castineira C, El Faedy O et al. The RIPASA score is sensitive and specific for the diagnosis of acute appendicitis in a Western population. *Int J Colorectal Dis.* 2017; 32(4): 491-7. doi: 10.1007/s00384-016-2713-4.
- Damani SAAR, Shah SSH, Hashami A, Mansoori MS. Effective diagnosis of acute appendicitis - comparison of RIPASA and Alvarado scoring systems. *J Surg Pakistan.* 2016; 21(3): 88-91.
- Shuaib A, Shuaib A, Fakhra Z, Marafi B, Alsharaf K, Behbehani A. Evaluation of modified Alvarado scoring system and RIPASA scoring system as diagnostic tools of acute appendicitis. *World J Emerg Med.* 2017; 8(4): 276-80. doi: 10.5847/wjem.j.1920-8642.2017.04.005.
- Özsoy Z, Yenidoğan E. Evaluation of the Alvarado scoring system in the management of acute appendicitis. *Turk J Surg.* 2017; 33(3): 200-4. doi: 10.5152/turkjsurg.2017.3539.
- Chisthi MM, Surendran A, Narayanan JT. RIPASA and air scoring systems are superior to Alvarado scoring in acute appendicitis: Diagnostic accuracy study. *Ann Med Surg (Lond).* 2020; 59: 138-42.

doi: 10.1016/j.amsu.2020.09.029.

15. Kothari D, Kothari A, Kalantri A, Bhambani P. Modified Alvarado scoring system as a diagnostic tool for acute appendicitis at a tertiary care teaching hospital, Central India: a cross-sectional study. *Int Surg J.* 2017; 4(8): 2439-44. doi: 10.18203/2349-2902.isj20173142
16. Pasumarthi V, Madhu CP. A comparative study of RIPASA score and ALVARADO score in diagnosis of acute appendicitis. *Int Surg J.* 2018; 5(3): 796-801.
17. Sanjive JG, Ramaiah RH. Comparison of RIPASA and Alvarado scoring in the diagnosis of acute appendicitis and validation of RIPASA scoring. *Int Surg J.* 2019; 6(3): 935-9.
18. Noor S, Wahab A, Afridi G, Ullah K. Comparing RIPASA score and Alvarado score in an accurate diagnosis of acute appendicitis *J Ayub Med Coll Abbottabad.* 2020; 32(1): 38-41.
19. Malik MU, Connelly TM, Awan F, Pretorius F, Fiuza-Castineira C, El Faedy O et al. The RIPASA score is sensitive and specific for the diagnosis of acute appendicitis in a Western population. *Int J Colorectal Dis.* 2017; 32(4): 491-7. doi: 10.1007/s00384-016-2713-4.
20. Karami MY, Niakan H, Zadebagheri N, Mardani P, Shayan Z, Deilami I. Which one is better? comparison of the acute inflammatory response, Raja Isteri Pengiran Anak Saleha appendicitis and Alvarado scoring systems. *Ann Coloproctol.* 2017; 33(6): 227-31. doi: 10.3393/ac.2017.33.6.227.
21. Korkut M, Bedel C, Karancı Y, Avci A, Duyan M. Accuracy of Alvarado, Eskelinen, Ohmann, RIPASA and Tzanakis scores in diagnosis of acute appendicitis; a cross-sectional study. *Arch Acad Emerg Med.* 2020; 8(1): e20.

